

Improving Cattle Fertility with Genetics

The fertility of a breeder herd has a significant impact on the profitability of a beef breeding business. Basically, more calves from a set number of breeders over a defined period (12 months) equals more dollars.

The next bull you purchase is one factor that will impact on the fertility of your breeder herd. This will arise from keeping a selection of his daughters to enter the breeding herd (i.e. Self replacing herd).

The question to ask yourself is "How do I know that the bull(s) I am selecting will have the genetic package to produce very fertile daughters?"

There are currently two pieces of information that can be used as a guide to selecting bulls with "good" fertility genetics. These being genetic differences (BREEDPLAN Estimated Breeding Values) for Scrotal Size (SS) and Days to Calving (DC).

SCROTAL SIZE

Scrotal Size (SS) EBVs are estimates of the genetic differences between animals in scrotal circumference (cm) at 400 days of age.

Scrotal size is measured by BREEDPLAN members from behind the animal (Figure 1). The scrotum containing both descended testes is measured using a tape positioned where scrotal circumference is greatest. At the same time, the scrotum can be palpated to detect abnormalities.

Increased SS is associated with increased semen production in bulls, and earlier age at puberty of bull and heifer progeny. SS also has a small favourable relationship with Days to Calving (DC), such that bulls with larger SS tend to have daughters with shorter DC.

Larger, more positive, SS EBVs are generally more favourable. For example, a bull with an SS EBV of +1.7 cm would be expected to produce sons with larger testicles at yearling age and daughters that reach puberty earlier than the progeny of a bull with an SS EBV of +0.4 cm.

FAST // FACTS

- More positive Scrotal Size EBVs (cm) are more favourable and associated with larger testicle size in yearling bulls and earlier age of puberty in heifers
- Lower or more negative Days to Calving EBVs (days) are more favourable and are associated with early heifer puberty, return to oestrus after calving and early conception

A raw SS (eg 36cm) and SS EBV (+1.7 cm) on sale bulls should not be confused as they indicate slightly different traits. A raw SS should be used to indicate if a bull has a higher probability of being fertile himself (i.e. is able to get his share of cows in calf). The threshold for 2 year old Bos indicus sale bulls on "good" nutrition is 34 cm.

Bulls selected for high SS EBVs are expected on average to breed daughters with earlier puberty and with shorter days to calving.



Figure 1. Measuring Scrotal Circumference

DAYS TO CALVING

Days to Calving (DC) EBVs are estimates of genetic differences between animals in female fertility, expressed as the number of days from the start of the joining period (ie. when the female is introduced to a bull) until subsequent calving (Figure 2).

DC promotes those cows that calve earlier in the season compared to those that calve later, while penalising those cows that do not calve. Negative DC EBVs indicate a shorter interval from the start of joining season until calving.

Variation in DC is mainly due to differences in the time taken for females to conceive after the commencement of the joining period (Figure 2). Females with shorter DC EBVs tend to be those which also show early puberty as heifers, which return to oestrous earlier after calving and conceive early in the joining period.

Days to calving EBVs may be reported for sires and dams - however it is likely that only animals with reasonable accuracies will appear in catalogues. The DC EBVs are reported in days and will generally have low accuracies until a sire has a large number of daughters with days to calving records.

Lower or negative DC EBVs are generally more favourable and indicate sires that will produce daughters with shorter number of days to calving and this effect will be cumulative over the life of those daughters. For example, a bull with a DC EBV of -10 days would be expected to produce daughters that conceive earlier in

the joining period than the daughters of a bull with a DC EBV of +10 days.

Note that there is only a small favourable relationship between SS and DC. Therefore it is strongly recommended that you select for the DC trait directly if you want to improve female fertility.

CONCLUSION

Progress can be made in fertility by selecting bulls that have the right genetic package. Indicators of bulls with a good fertility genetic package are those with an above average SS EBV and below average (negative) DC EBV.

You will notice in many sale catalogues that have weight EBVs available for selection do not have fertility EBVs available (eg SS & DC). We encourage all BREEDPLAN members to record these fertility traits, submit for analyses and have EBVs available for their own selection and for their clients.

Commercial bull buyers should ask for this information on the bulls they will be potentially purchasing and taking home. Other EBVs (weight, carcass), vet inspections and visual appraisal should also be included in your bull buying decisions.

For further information on the fertility EBVs or for stud breeders who wish to record the information to generate them, contact:

Paul Williams, Tropical Beef Technology Services
T: 07 4927 6066 or E: paul@sbts.une.edu.au




Figure 2. Measure of Days to Calving